

Nutrition and post-ICU recovery

Reference list

1. Iwashyna TJ, Ely EW, Smith DM, Langa KM. Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA*. 2010;304(16):1787-1794.
2. Herridge MS, Tansey CM, Matte A, et al. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med*. 2011;364(14):1293-1304.
3. Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin Nutr*. 2006;84(3):475-482.
4. Puthuchery ZA, Rawal J, McPhail M, et al. Acute skeletal muscle wasting in critical illness. *JAMA*. 2013;310(15):1591-1600.
5. Parry SM, El-Ansary D, Cartwright MS, et al. Ultrasonography in the intensive care setting can be used to detect changes in the quality and quantity of muscle and is related to muscle strength and function. *J Crit Care*. 2015;30(5):1151 e1159-1114.
6. Mayer KP, Thompson Bastin ML, Montgomery-Yates AA, et al. Acute skeletal muscle wasting and dysfunction predict physical disability at hospital discharge in patients with critical illness. *Crit Care*. 2020;24(1):637.
7. van Gassel RJJ, Baggerman MR, van de Poll MCG. Metabolic aspects of muscle wasting during critical illness. *Curr Opin Clin Nutr Metab Care*. 2020;23(2):96-101.
8. Sherwood L. *Human Physiology: From Cells to Systems*. Vol 9th Ed.2016.
9. Ridley EJ, Peake SL, Jarvis M, et al. Nutrition Therapy in Australia and New Zealand Intensive Care Units: An International Comparison Study. *JPEN*. 2018;42(8):1349-1357.
10. Ridley EJ, Chapple LS, Chapman MJ. Nutrition intake in the post-ICU hospitalization period. *Curr Opin Clin Nutr Metab Care*. 2020;23(2):111-115.
11. Ferrie S, Allman-Farinelli M, Daley M, Smith K. Protein Requirements in the Critically Ill: A Randomized Controlled Trial Using Parenteral Nutrition. *JPEN*. 2016;40(6):795-805.
12. Fetterplace K, Deane AM, Tierney A, et al. Targeted Full Energy and Protein Delivery in Critically Ill Patients: A Pilot Randomized Controlled Trial (FEED Trial). *JPEN*. 2018.
13. McNelly AS, Bear DE, Connolly BA, et al. Effect of Intermittent or Continuous Feed on Muscle Wasting in Critical Illness: A Phase 2 Clinical Trial. *Chest*. 2020;158(1):183-194.
14. Morton RW, Traylor DA, Weijs PJM, Phillips SM. Defining anabolic resistance: implications for delivery of clinical care nutrition. *Curr Opin Crit Care*. 2018;24(2):124-130.
15. Wall BT, Hamer HM, de Lange A, et al. Leucine co-ingestion improves post-prandial muscle protein accretion in elderly men. *Clin Nutr*. 2013;32(3):412-419.
16. Leenders M, Verdijk LB, van der Hoeven L, et al. Prolonged leucine supplementation does not augment muscle mass or affect glycemic control in elderly type 2 diabetic men. *J Nutr*. 2011;141(6):1070-1076.
17. Verhoeven S, Vanschoonbeek K, Verdijk LB, et al. Long-term leucine supplementation does not increase muscle mass or strength in healthy elderly men. *Am J Clin Nutr*. 2009;89(5):1468-1475.
18. Bear DE, Langan A, Dimidi E, et al. beta-Hydroxy-beta-methylbutyrate and its impact on skeletal muscle mass and physical function in clinical practice: a systematic review and meta-analysis. *Am J Clin Nutr*. 2019;109(4):1119-1132.